

What is the Correct $\pi^-p \rightarrow \omega n$ Cross Section at Threshold?[†]

G. Penner* and U. Mosel

Institut für Theoretische Physik, Universität Giessen, D-35392 Giessen

The $\pi^-p \rightarrow \omega n$ threshold cross section of refs. [1, 2, 3] resisted up to now a consistent theoretical description in line with experiment, mainly caused by too large Born contributions. This led to a discussion in the literature about the correctness of the extraction of these data points. We show that the extraction method used in these references is indeed correct and that there is no reason to doubt the correctness of these data.

PACS numbers: 13.75.Gx

I. INTRODUCTION

In the 70's a series of experiments [1, 2, 3] was performed to measure the $\pi^-p \rightarrow \omega n$ cross section just above threshold. These data resisted up to now a consistent theoretical description, mainly caused by too large Born contributions [4]. As a consequence these diagrams were either neglected [5, 6] or suppressed by very soft formfactors [9]. These findings motivated a discussion in the literature about the experimentalists' way to extract the two-body cross section [7] and readjustments of the published $\pi^-p \rightarrow \omega n$ cross section data were performed [8, 9, 10].

The cause of the discussion is the experimentalists' unusual method to cover the full range of the ω spectral function. An integration over at least one kinematical variable is necessary to make sure that all pion triples with invariant masses around m_ω are taken into account, so that the ω spectral function with a width of 8 MeV is well covered. Instead of fixing the incoming pion momentum and integrating out the invariant mass of the pion triples directly, the authors of [1, 2, 3] fixed the outgoing neutron laboratory momentum and angle and performed an integration over the incoming pion momentum.

Led by the observation that the cross sections of [1, 2, 3] result in a hard-to-understand energy dependence of the transition matrix element the authors of [7] claimed that due to the experimental method just described the count rates covered only a fraction of the ω spectral function. As a consequence, the authors of [7, 8, 9] advocated that the two-body total cross sections given in [3, 11] should be modified. Imposing this modification of the threshold cross section, in [7] a practically constant transition matrix element up to 1.74 GeV corresponding to an S_{11} or D_{13} [12] (since the pion momentum is almost constant in this region) wave production mechanism (in the usual πN notation, i.e. an $IJ^P = \frac{1}{2}\frac{1}{2}^-$ or $IJ^P = \frac{1}{2}\frac{3}{2}^-$, resp., partial wave) was deduced.

This modification, however, immediately raises another problem. Taking into account this “spectral func-




FIG. 1: S_{11} inelastic partial wave cross section of $\pi N \rightarrow \pi N$ as deduced from SM00 [13] $\circ$ and $\pi N \rightarrow 2\pi N$ partial wave cross section as deduced by [14] $\bullet$. The curve gives the result of a coupled-channel analysis [15].

tion correction” increases the ωN cross section for $1.72 \leq \sqrt{s} \leq 1.74$ up to $\sigma \geq 3$ mb, which is in contradiction to the inelasticity deduced from $\pi N \rightarrow \pi N$ partial wave analyses (e.g. SM00 [13]). As can be seen in fig. 1 the $\pi N \rightarrow \pi N$ inelasticity in the S_{11} partial wave is already saturated by $\pi N \rightarrow 2\pi N$ in this energy region, i.e. an additional ωN contribution of $\sigma^{S_{11}} \geq 3$ mb would lead to a gross overestimate of the inelasticity:

$$\begin{aligned} \sigma_{in}^{S_{11}} &\geq \sigma_{2\pi N}^{S_{11}} + \sigma_{\omega N}^{S_{11}} \\ &\geq 2.5 \text{ mb} + 3 \text{ mb} \gg 2.4 \text{ mb} \approx \sigma_{in}^{S_{11}} (\text{SM00}) . \end{aligned} \quad (1)$$

The same argument holds for the D_{13} wave (see fig. 5 in [15]).

Because of this inconsistency of the newly extracted cross sections with existing inelasticities and because of the importance of the $\pi N \rightarrow \omega N$ cross section for both unitary models analyzing reactions on the nucleon in the c.m. energy range $1.7 \leq \sqrt{s} \leq 2.0$ GeV [15] and in-medium models of vector mesons [5, 16, 17] we reanalyze the extraction method used in refs. [1, 2, 3] by presenting a complete derivation (given in parts in [1]) of the relation between the experimental count rates and the extracted two-body cross section for $\pi N \rightarrow \omega N$.

[†]Work supported by DFG and GSI Darmstadt.

*Electronic address: gregor.penner@theo.physik.uni-giessen.de

II. COUNT RATES AND TWO-BODY CROSS SECTIONS

The experimental count rates are given by (B3) (we label all equations to be found in [1] with the letter B and those in [7] by the letter H):

$$\bar{N} = N_H \int_{\Delta\Omega_L} \int_{\Delta p'_L} \frac{d^3\sigma}{d\Omega_L dp'_L} d\Omega_L dp'_L, \quad (2)$$

where the cross section σ describes the process $\pi^- p \rightarrow n\pi^+\pi^-\pi^0$. All variables are taken in the c.m. system, unless they are denoted by the label L (laboratory frame). p (k) and p' (q) denote the incoming proton and outgoing neutron (incoming π and outgoing ω) four momenta. Absolute values of three momenta are denoted by upright letters, s and t are the usual Mandelstam variables. Ω_L is the neutron laboratory solid angle and N_H the number of target particles per unit area. The integral ranges in (2) refer to the binning of the count rates, i.e. they are the integrals to be performed for averaging over the experimental resolution intervals $\Omega_L \pm \Delta\Omega_L$ and $p'_L \pm \Delta p'_L$ and are not related to an integration over the ω spectral function. This will become clearer below (cf. eqn. (12)). The kinematics of the reaction are extracted from the center values of these intervals. Using

$$\begin{aligned} t = (p_L - p'_L)^2 &= m_p^2 + m_n^2 - 2m_p E_{nL} \\ &= m_p^2 + m_n^2 - 2m_p \sqrt{m_n^2 + p'_L{}^2} \end{aligned} \quad (3)$$

and

$$\begin{aligned} (q_L - p_L)^2 &= (k_L - p'_L)^2, \quad E_{\omega L} = E_{\pi L} + m_p - E_{nL} \\ \Rightarrow q^2 &= m_\pi^2 + m_n^2 + m_p^2 + \\ &\quad 2(m_p E_{\pi L} - m_p E_{nL} - E_{\pi L} E_{nL} + k_L p'_L x_L) \end{aligned} \quad (4)$$

($x_L = \cos \vartheta_L$) one finds

$$\bar{N} = \frac{N_H}{2\pi} \int_{\Delta\Omega_L} \int_{\Delta p'_L} \frac{d^2\sigma}{dt d\sqrt{q^2}} J_P d\Omega_L dp'_L,$$

assuming the cross section is independent of the neutron azimuthal angle φ . The Jacobian J_P is given by

$$J_P = \left| \frac{d\sqrt{q^2}/dp'_L}{dt/dp'_L} \frac{d\sqrt{q^2}/dx_L}{dt/dx_L} \right| = \frac{2m_p k_L p'_L{}^2}{\sqrt{q^2} E_{nL}},$$

because $dt/dx_L = 0$. Since in the actual experiment the time of flight τ_L of the neutron over a distance d ,

$$\tau_L = \frac{1}{\beta_L} \frac{d}{c}$$

with the velocity $\beta_L = p'_L/E_{nL}$, is measured – not its three-momentum –, the count rate is reexpressed in terms of the time of flight:

$$\bar{N} = \frac{N_H}{2\pi} \int_{\Delta\Omega_L} \int_{\Delta\tau_L} \frac{d^2\sigma}{dt d\sqrt{q^2}} J_\tau d\Omega_L d\tau_L,$$

with

$$J_\tau = J_P \frac{dp'_L}{d\tau_L}.$$

The factor linking the Jacobians is

$$\frac{dp'_L}{d\tau_L} = \frac{dp'_L}{d\beta_L} \frac{d\beta_L}{d\tau_L} = \frac{p'_L{}^2 E_{nL}}{m_n^2} \frac{c}{d} \text{ because of } \frac{d\beta_L}{dp'_L} = \frac{m_n^2}{E_{nL}^3}.$$

We now relate this count rate to the two-body cross section [18]

$$\frac{d\sigma^{2b}}{dt} = \frac{1}{64\pi s k^2} \frac{1}{2} \sum_{\lambda_p, \lambda_n, \lambda_\omega} |\mathcal{M}(s, t)|^2 \quad (5)$$

of $\pi^- p \rightarrow \omega n$, assuming a stable ω . In order to do so, we deviate from the derivation in [1] and start with the general cross section formula for $\pi N \rightarrow 3 + 4 + 5 + \dots + l$ in the c.m. system [18]:

$$\begin{aligned} d\sigma &= \frac{1}{4k\sqrt{s}} \frac{1}{2} \sum |\tilde{\mathcal{M}}|^2 (2\pi)^4 \prod_{j=3}^l \frac{d^3 k'_j}{(2\pi)^3 2E'_j} \\ &\quad \times \delta^4(p + k - \sum_{j=3}^l k'_j), \end{aligned} \quad (6)$$

where the sum stands for summing over initial and final spins. Here (for simplicity, we assume that the ω only decays into 3 pions: $\pi^- p \rightarrow n\omega \rightarrow n\pi^+\pi^-\pi^0$), the matrix element reads:

$$\tilde{\mathcal{M}} = \mathcal{M}_{\pi^- p \rightarrow \omega n}^\mu D_{\mu\nu}^\omega(q^2) \mathcal{H}_{\omega \rightarrow 3\pi}^\nu$$

with $D_{\mu\nu}^\omega(q^2) = \sum_{\lambda_\omega} \varepsilon_\mu^\dagger(\lambda_\omega) \varepsilon_\nu(\lambda_\omega) \Delta_\omega(q^2)$ and $\Delta_\omega(q^2) = (q^2 - m_\omega^2 + i\sqrt{q^2} \Gamma_{\omega \rightarrow 3\pi})^{-1}$. Thus, in $|\tilde{\mathcal{M}}|^2$ a sum over λ_ω and λ'_ω appears. However, since the ω decay amplitude $\varepsilon_{\lambda_\omega} \cdot \mathcal{H}_{\omega \rightarrow 3\pi}$ can be decomposed into spherical harmonics $Y_{1\lambda_\omega}$ and the outgoing pion angles are integrated out, there are only contributions for $\lambda_\omega = \lambda'_\omega$. To introduce the ω spectral function $\rho_\omega(q^2)$ in (6), we note that we can evaluate the decay $\varepsilon_{\lambda_\omega} \cdot \mathcal{H}_{\omega \rightarrow 3\pi}$ in the ω rest frame, which is hence independent of the polarization λ_ω . Then the width of the ω is given by

$$\begin{aligned} \Gamma_{\omega \rightarrow 3\pi}(q^2) &= \frac{1}{2\sqrt{q^2}} \int \prod_{j=4}^6 \frac{d^3 k'_j}{(2\pi)^3 2E'_j} |\varepsilon_{\lambda_\omega}^\mu \cdot \mathcal{H}_\mu|^2 \\ &\quad \times (2\pi)^4 \delta^4(q - \sum_{j=4}^6 k'_j), \end{aligned} \quad (7)$$

valid for any λ_ω , and related to the ω spectral function in the following way:

$$\begin{aligned} \rho_\omega(q^2) &= -\frac{1}{\pi} \text{Im} \Delta_\omega(q^2) \\ &= \frac{1}{\pi} |\Delta_\omega(q^2)|^2 \sqrt{q^2} \Gamma_\omega(q^2) \\ &= \frac{1}{\pi} |\Delta_\omega(q^2)|^2 \frac{1}{2} \int \prod_{j=4}^6 \frac{d^3 k'_j}{(2\pi)^3 2E'_j} |\varepsilon_{\lambda_\omega}^\mu \cdot \mathcal{H}_\mu|^2 \\ &\quad \times (2\pi)^4 \delta^4(q - \sum_{j=3}^6 k'_j). \end{aligned}$$

Now, we can rewrite the cross section of eqn. (6) by introducing $1 = \int d^4q \delta^4(q - \sum_{j=3}^6 k'_j)$ and using the spectral

function:

$$\begin{aligned}
d^2\sigma &= \frac{1}{4k\sqrt{s}} \frac{1}{2} \sum_{\lambda_p, \lambda_n, \lambda_\omega} \int |\mathcal{M} \cdot \varepsilon_{\lambda_\omega}|^2 (2\pi)^4 \delta^4(p + k - p' - q) \frac{d^3p'}{(2\pi)^3 2E_n} \frac{2\pi}{(2\pi)^4} \rho_\omega(q^2) d^4q \\
&= \frac{1}{4k\sqrt{s}} \frac{1}{(2\pi)^2} \frac{1}{2} \sum_{\lambda_p, \lambda_n, \lambda_\omega} \int |\mathcal{M} \cdot \varepsilon_{\lambda_\omega}|^2 \frac{d^3p'}{2E_n} \rho_\omega(q^2) \Bigg|_{q=p+k-p'} \\
&= \frac{1}{8k\sqrt{s}} \frac{1}{2\pi} \frac{1}{2} \sum_{\lambda_p, \lambda_n, \lambda_\omega} |\mathcal{M} \cdot \varepsilon_{\lambda_\omega}|^2 \frac{p'^2 dp'}{E_n} dx \rho_\omega(q^2) \Bigg|_{q=p+k-p'} \\
&= \frac{p'}{32\pi\sqrt{s}k^2} \frac{1}{2} \sum_{\lambda_p, \lambda_n, \lambda_\omega} |\mathcal{M} \cdot \varepsilon_{\lambda_\omega}|^2 \frac{dp' dt}{E_n} \rho_\omega(q^2) \Bigg|_{q=p+k-p'} \\
&\stackrel{(5)}{=} 2\sqrt{q^2} \frac{d\sigma^{2b}}{dt} \rho_\omega(q^2) dt d\sqrt{q^2} \Bigg|_{q=p+k-p'} ,
\end{aligned} \tag{8}$$

where we have used

$$\begin{aligned}
t = (p - p')^2 &= m_p^2 + m_n^2 - 2(E_p E_n - k p' x) \\
\Rightarrow \left| \frac{dt}{dx} \right| &= 2k p' ,
\end{aligned} \tag{9}$$

and

$$\begin{aligned}
E_n &= \frac{s + m_n^2 - q^2}{2\sqrt{s}} \\
\Rightarrow \frac{dp'}{d\sqrt{q^2}} &= \frac{dp'}{dE_n} \frac{dE_n}{d\sqrt{q^2}} = \frac{\sqrt{q^2} E_n}{p' \sqrt{s}} .
\end{aligned} \tag{10}$$

Thus we have

$$\frac{d^2\sigma}{dt d\sqrt{q^2}} = 2\sqrt{q^2} \frac{d\sigma^{2b}}{dt} \rho_\omega(q^2) \Bigg|_{q=p+k-p'}$$

and finally for the experimental count rate

$$\bar{N} = \frac{N_H}{2\pi} \int_{\Delta\Omega_L} \int_{\Delta\tau_L} 2\sqrt{q^2} \frac{d\sigma^{2b}}{dt} \rho_\omega(q^2) \Bigg|_{q=p+k-p'} J_\tau d\Omega_L d\tau_L . \tag{11}$$

To eliminate the ω spectral function, the experimentalists [1, 2, 3] performed an integration over the incoming pion three-momentum k_L by summing over all beam settings, which can be transformed into an integration over the ω four-momentum squared:

$$\begin{aligned}
N &= \int dk_L \bar{N} \\
&= \frac{N_H}{2\pi} \int dk_L \int_{\Delta\Omega_L} \int_{\Delta\tau_L} 2\sqrt{q^2} \frac{d\sigma^{2b}}{dt} \rho_\omega(q^2) J_\tau d\Omega_L d\tau_L
\end{aligned}$$

$$= \frac{N_H}{2\pi} \int_{\Delta\Omega_L} \int_{\Delta\tau_L} \frac{d\sigma^{2b}}{dt} J_k J_\tau d\Omega_L d\tau_L , \tag{12}$$

where we have used the normalization of the spectral function $\int \rho_\omega(q^2) dq^2 = 1$ and the assumption that the matrix element in eqn. (5) only varies slightly around the peak of the ω spectral function¹ for fixed t corresponding to fixed p'_L via eqn. (3)². In the first line, due to four-momentum conservation, the ω momentum is restricted to $q = p + k - p'$. The Jacobian J_k is given by

$$J_k \equiv \frac{dk_L}{d\sqrt{q^2}} \stackrel{(4)}{=} \frac{\sqrt{q^2}}{\frac{k_L}{E_{\pi L}} (m_p - E_{nL}) + p'_L x_L} .$$

In the derivation of eqn. (12) two more assumptions enter that are checked in the following:

- Sufficient coverage of the ω spectral function for all kinematics extracted, even at low p' values:
The incoming pion lab momentum range was $k_L \in [1.04, 1.265]$ GeV, which translates into a c.m. energy range of $\sqrt{s} \in [1.6938, 1.8133]$ GeV. Using $p' \in [0.03, 0.21]$ GeV, this leads to the following ranges for the upper and lower limits in the q^2 integration: $q_+^2 \in [0.822^2, 0.869^2]$ GeV² and $q_-^2 \in [0.7^2, 0.753^2]$ GeV². Even in the worst case

¹ The approximate constancy of the matrix element is the basic assumption for extracting the two-body cross section from **any** experiment dealing with decaying final state particles.

² Remember that the integral range $\Delta\tau_L$ in eqn. (12) corresponds to the experimental resolution.




FIG. 2: Differential cross section for backward ω c.m. angles. The data points are from $\bullet$: [1], $\triangle$: [2], $\circ$: [3], and $\square$: [19]. The curve gives the result of a coupled-channel analysis [15].

the integration extends over at least 7 half widths $\Gamma_\omega/2$ on either side of m_ω^2 and thus covers more than 92% of the spectral function.

- Constancy of the product of the Jacobians $J_k J_\tau$: By using eqns. (3), (4), (9), and (10) one can easily show that the product

$$J_k J_\tau = \frac{2m_p k_L p_L'^4}{m_n^2 \left[\frac{k_L}{E_{\pi_L}} (m_p - E_{n_L}) + p_L' x_L \right]} \frac{c}{d}$$

varies for fixed t by less than 2.5 percent in the interval $\sqrt{q^2} \in [m_\omega - 2\Gamma_\omega, m_\omega + 2\Gamma_\omega]$ for all kinematics considered in the experiment.

Since the countrate of eqn. (12) is identical to the one given between eqns. (B9) and (B10) in [1], the method used in refs. [1, 2, 3] and correspondingly their two-body cross sections are correct.

It is important to notice that in the data analysis both p_L' and x_L are needed to fix the kinematics of the measured events. During the count rate corrections (flux normalization in dependence on k_L , background subtraction via missing mass spectra), for each k_L beam setting the measured p_L' and x_L translate into $\sqrt{q^2}$ (see (4)) and can also be Lorentz transformed into their c.m. values $p' = p'(E_{\pi_L}, p_L', x_L)$ and $x = x(E_{\pi_L}, p_L', x_L)$. The events can now be regrouped in p' , $\sqrt{q^2}$, and x intervals. Then, after having performed the integration over all $\sqrt{q^2}$ as in (12), the translation of a given p' into $\sqrt{s}$ can only be done by assuming that the main contribution to the corrected count rates comes from around the peak of the omega spectral function $q^2 \sim m_\omega^2$.

The main assertion of ref. [7], manifested in eqn. (H4), is that instead of (12) only a fraction of the cross section for the production of an unstable particle had been measured in [1, 2, 3]. This fraction is determined by translating the experimental p' binning intervals given in [3] into interval bounds for the integration over the ω spectral function. Equation (H4), which is used for the cross




FIG. 3: Differential cross section for forward ω c.m. angles. For the notations see fig. 2.

section corrections in [8, 9, 10], is identical to the third line of eqn. (8) under the assumption that p' is bound to the p' binning intervals. However, as pointed out above, in an experimental event E_{π_L} , p_L' , and x_L are fixed and hence eqn. (11) has to be applied to the experimental count rate for a fixed pion momentum.

The experimental integration over the incoming pion momentum is introduced in ref. [7] only in the subsequent discussion between eqns. (H9) and (H10). In the paragraph following eqn. (H16) the authors of ref. [7] argue that the range of this integral is narrowed due to the p' binning (as in (H4)). The ω mass is thus allowed to vary only in the interval given by the p' interval ranges for a fixed pion momentum. But as shown above, fixing p' only fixes the incoming pion momentum if one assumes a specific ω mass. Hence the pion momentum integration performed in the data analysis indeed translates into an ω mass integration only bounded by the pion momentum range and thus leads to eqn. (12). This relation between the neutron momentum p' , the pion momentum k , and the ω mass were thus treated improperly in ref. [7].

The second correction factor extracted in ref. [7] due to the neutron momentum binning of $\Delta p' = 10$ MeV is nothing but the result of averaging the third line of eqn. (8) over the neutron c.m. momentum: $(\Delta p')^{-1} \int_{p'-\Delta p'}^{p'+\Delta p'} \tilde{p}^2 d\tilde{p} = p'^2 + (\Delta p')^2/12$ (cf. (H10)). This differs at most (“worst” case: $p' = 30$ MeV) by 1 percent from p'^2 and is therefore negligible.

Furthermore, it is obvious from figs. 2 and 3 that the differential ωN data from all three references [1, 2, 3]³ are completely in line with each other and also with ref. [19]⁴. The same holds true for the total cross sections of

³ The total cross sections given in refs. [1, 2] are actually angle-differential cross sections (mostly at forward and backward neutron c.m. angles) multiplied with 4π .

⁴ The differential cross sections are extracted from the corrected cosine event distributions given in ref. [19] with the help of their




FIG. 4: $\pi N \rightarrow \omega N$ total cross section. Data are from $\circ$: [3], ∇ : [11], $\square$: [19]. The curve gives the result of a coupled-channel analysis [15].

ref. [3] in comparison with other experiments⁵, see fig. 4. There is, therefore, no reason to hypothesize – as in [10] – that the formalism developed in [1] could have been used incorrectly in [2] and [3].

total cross sections.

⁵ Note, that all other experiments measured $\pi^+ n \rightarrow \omega p$.

In this context we stress one more point. Very close to threshold, the two-body cross section $\pi^- p \rightarrow \omega n$ extracted from experimental count rates could be influenced by the strong πN interaction for slow pions stemming from $\omega \rightarrow 3\pi$. However, this point was checked in ref. [2] by also looking at $\omega \rightarrow \pi^0 \gamma$; they did not find any deviations between the two ways of extraction.

III. SUMMARY AND CONCLUSION

We have shown that the extraction method presented in [1] and also used in [2, 3] is indeed correct. There is no reason to doubt the correctness of the data presented in these references; they are in line with each other and also with other experimental data. The reanalysis of the ωN -production data in ref. [7], on which the theoretical descriptions of refs. [8, 9] are based, as well as the speculations in ref. [10] thus lack any basis. In ref. [15] we show how the $\pi^- p \rightarrow \omega n$ cross section at threshold can be understood in a coupled-channel analysis.

-
- [1] D.M. Binnie et al., Phys. Rev. D **8**, 2789 (1973).
 - [2] J. Keyne, D.M. Binnie, J. Carr, N.C. Debenham, A. Duane, D.A. Garbutt, W.G. Jones, I. Siotis, and J.G. McEwen, Phys. Rev. D **14**, 28 (1976).
 - [3] H. Karami, J. Carr, N.C. Debenham, D.A. Garbutt, W.G. Jones, D.M. Binnie, J. Keyne, P. Moissidis, H.N. Sarma, and I. Siotis, Nucl. Phys. **B154**, 503 (1979).
 - [4] F. Klingl, Ph.D. thesis, University of Munich, Hieronymus, Munich 1998.
 - [5] M. Post and U. Mosel, Nucl. Phys. **A688**, 808 (2001).
 - [6] M. Lutz, G. Wolf, and B. Friman, Nucl. Phys. **A661**, 526c (1999) and *Proceedings of the International Workshop XXVIII on Gross Properties of Nuclei and Nuclear Excitations: Hirschegg, Austria, 16-22 Jan. 2000*, Editors: M. Buballa, B.-J. Schaefer, W. Nörenberg, J. Wambach, GSI; e-Print ArXiv: [nucl-th/0003012](#).
 - [7] C. Hanhart and A. Kudryavtsev, Eur. Phys. J. A **6**, 325 (1999).
 - [8] A. Sibirtsev and A. Cassing, Eur. Phys. J. A **7**, 407 (2000).
 - [9] A.I. Titov, B. Kämpfer, and B.L. Reznik, e-Print ArXiv: [nucl-th/0102032](#).
 - [10] C. Hanhart, A. Sibirtsev, and J. Speth, e-Print ArXiv: [nucl-th/0107245](#).
 - [11] H. Höhler, *Landolt-Börnstein*, Vol. 9, Springer, Berlin 1983.
 - [12] We thank C. Hanhart for pointing out this second possibility to us.
 - [13] M.M. Pavan, R.A. Arndt, I.I. Strakovsky, and R.L. Workman, Phys. Scripta **T87** 62 (2000), e-Print ArXiv: [nucl-th/9807087](#), R.A. Arndt, I.I. Strakovsky, R.L. Workman, and M.M. Pavan, Phys. Rev. C **52**, 2120 (1995), updates available via: <http://gwdac.phys.gwu.edu/>.
 - [14] D.M. Manley, R.A. Arndt, Y. Goradia, and V.L. Teplitz, Phys. Rev. D **30**, 904 (1984).
 - [15] G. Penner and U. Mosel, submitted to Phys. Rev. C, e-Print ArXiv: [nucl-th/0111023](#).
 - [16] B. Friman, *Proceedings of the APCTP-Workshop on Astro-Hadron Physics, Seoul, Korea, October 1997*, e-Print ArXiv: [nucl-th/9801053](#).
 - [17] F. Klingl, T. Waas, and W. Weise, Nucl. Phys. **A650**, 299 (1999).
 - [18] C. Itzykson and J.-B. Zuber, *Quantum Field Theory*, McGraw-Hill, New York 1980.
 - [19] J.S. Danburg, M.A. Abolins, O.I. Dahl, D.W. Davies, P.L. Hoch, J. Kirz, D.H. Miller, and R.K. Rader, Phys. Rev. D **2**, 2564 (1970).